

# GALOIS SYMMETRIES OF KNOT SPACES

PEDRO BOAVIDA DE BRITO AND GEOFFROY HOREL

ABSTRACT. We exploit the Galois symmetries of the little disks operads to show that many differentials in the Goodwillie-Weiss spectral sequences approximating the homology and homotopy of knot spaces vanish at a prime  $p$ . Combined with recent results on the relationship between embedding calculus and finite-type theory, we deduce that the  $(n + 1)$ -st Goodwillie-Weiss approximation is a  $p$ -local universal Vassiliev invariant of degree  $\leq n$  for every  $n \leq p + 1$ .

A *knot* is a smooth embedding  $\mathbb{R} \rightarrow \mathbb{R}^3$  which coincides with the standard inclusion  $t \mapsto (t, 0, 0)$  in the complement of  $[0, 1]$ . The space of such, with the weak Whitney topology, is denoted  $K$ . In this paper, we show that the Goodwillie-Weiss approximation to the space of knots gives a  $p$ -local analog of Kontsevich’s integral, in a range which improves as the prime  $p$  increases.

**Theorem A.** *Let  $p$  be a prime and  $n \leq p + 1$ . The  $(n + 1)$ -st Goodwillie-Weiss approximation of the space of knots*

$$K \rightarrow T_{n+1}(K)$$

*is, on path components, a universal finite type invariant of degree  $n$  over  $\mathbb{Z}_{(p)}$ , the localization of  $\mathbb{Z}$  at the prime  $p$ .*

*Furthermore, under the same assumption on  $n$ , there is a non-canonical  $p$ -local isomorphism (i.e. an isomorphism after tensoring with  $\mathbb{Z}_{(p)}$ )*

$$\pi_0(T_{n+1}(K)) \cong \bigoplus_{s \leq n} \mathcal{A}_s^I$$

*where  $\mathcal{A}_s^I$  is the algebra of indecomposable Feynman diagrams, that is, the quotient of the free abelian group on univalent trees of degree  $s$  by the antisymmetry, IHX and  $STU^2$  relations.*

We now describe in more detail the main characters of this story: the *Vassiliev tower* and the *Goodwillie-Weiss tower*.

A finite-type invariant of degree  $n$  is a homomorphism from  $\pi_0(K)$  (viewed as a monoid with respect to concatenation) to an abelian group, whose extension<sup>1</sup> to singular knots having  $n + 1$  double points vanishes. It is not known whether finite type invariants – also called Vassiliev invariants – distinguish all knots, but it is known that they abound. Examples include all so-called quantum invariants, and

---

We gratefully acknowledge the support through: grant SFRH/BPD/99841/2014 and project MAT-PUR/31089/2017, funded by Fundação para a Ciência e Tecnologia; projects ANR-14-CE25-0008 SAT, ANR-16-CE40-0003 ChroK, ANR-18-CE40-0017 PerGAMo, funded by Agence Nationale pour la Recherche.

<sup>1</sup>Following Vassiliev, Goussarov and Birman-Lin, the extension  $\bar{\mu}$  of a knot invariant  $\mu$  to singular knots  $K$  having at most  $k$  double points is defined inductively by the formula:  $\bar{\mu}(K) = \bar{\mu}(K_+) - \bar{\mu}(K_-)$  where  $K_+$  and  $K_-$  denote the knots obtained from  $K$  by resolving a given double point in the two possible ways. Some standard references are [27], [2].

so all invariants coming from perturbative Chern-Simons theory, as well as many classically studied knot invariants. Two knots are declared to be  $n$ -equivalent if all finite type invariants of degree  $\leq n$  agree on these knots; this defines an equivalence relation  $\sim_n$  on  $\pi_0(K)$ . The equivalence relation  $\sim_n$  is finer than  $\sim_{n-1}$ , and therefore we have a tower of surjections

$$(0.1) \quad \pi_0(K) \rightarrow \dots \rightarrow \pi_0(K)/\sim_n \rightarrow \pi_0(K)/\sim_{n-1} \rightarrow \dots \rightarrow \pi_0(K)/\sim_1$$

that we call the *Vassiliev tower*. It turns out [17, Theorem 3.1] that the concatenation of knots endows each of the sets  $\pi_0(K)/\sim_n$  with the structure of a finitely generated abelian group. There is moreover a surjective map

$$\mathcal{A}_n^I \rightarrow \ker[\pi_0(K)/\sim_n \rightarrow \pi_0(K)/\sim_{n-1}] .$$

A deep and fundamental result in the development of the subject was Kontsevich's construction of the so-called Kontsevich integral which produces a rational inverse of this map and so a (non-canonical) isomorphism

$$(\pi_0(K)/\sim_n) \otimes \mathbb{Q} \cong \oplus_{s \leq n} \mathcal{A}_s^I \otimes \mathbb{Q} .$$

The *Goodwillie-Weiss tower*, on the other hand, is a tower of topological spaces

$$K \rightarrow \dots \rightarrow T_n(K) \rightarrow T_{n-1}(K) \rightarrow \dots \rightarrow T_2(K)$$

The map  $K \rightarrow T_n(K)$  records the value of an embedding  $\mathbb{R} \rightarrow \mathbb{R}^3$  on subsets of  $\mathbb{R}$  given by the complement of at most  $n+1$  points of  $(0, 1)$  and is, in a homotopical sense, universal with respect to this data. One can hit the Goodwillie-Weiss tower with  $\pi_0$  and obtain a tower

$$(0.2) \quad \pi_0(K) \rightarrow \dots \rightarrow \pi_0(T_n(K)) \rightarrow \pi_0(T_{n-1}(K)) \rightarrow \dots \rightarrow \pi_0(T_2(K)) .$$

It is known that all the sets  $\pi_0(T_n(K))$  are finitely generated abelian groups and that all the maps in the tower are compatible with this structure [8], [4]. There is a spectral sequence that computes  $\pi_0(T_n(K))$  (and also its higher homotopy groups):

$$E_{s,t}^1 \implies \pi_{t-s} T_n(K) .$$

The  $E^1$ -page consists of the homotopy groups of the homotopy fibers of the maps  $T_s(K) \rightarrow T_{s-1}(K)$  with  $s \leq n$  and can be expressed in terms of homotopy groups of spheres. The  $d^1$ -differential can be described explicitly and we have an isomorphism

$$E_{s,s}^2 \cong \mathcal{A}_{s-1}^I$$

(see [10] or [24] for an alternative account).

A long-standing conjecture is that the towers (0.1) and (0.2) are isomorphic; see the last paragraph of [15] and [9, Conjecture 1.1]. Some important steps have been made in this direction. Volić thesis [28] provided the first indication, though in the homological context. In [8], Budney-Conant-Koytcheff-Sinha showed that the map  $\pi_0(K) \rightarrow \pi_0(T_n(K))$  is a degree  $(n-1)$  invariant. A new proof of this result, using grope cobordisms, will appear in [21]. The upshot is that the tower (0.1) maps to the tower (0.2) (with a shift of degree), that is, we have a commutative ladder of abelian groups

$$\begin{array}{ccccccc} \dots & \longrightarrow & \pi_0(K)/\sim_n & \longrightarrow & \pi_0(K)/\sim_{n-1} & \longrightarrow & \dots \longrightarrow \pi_0(K)/\sim_1 \\ & & \downarrow & & \downarrow & & \downarrow \\ \dots & \longrightarrow & \pi_0(T_{n+1}(K)) & \longrightarrow & \pi_0(T_n(K)) & \longrightarrow & \dots \longrightarrow \pi_0(T_2(K)) . \end{array}$$

By work of Conant, Kosanović, Teichner and Shi [11, 20, 21], a sufficient condition for the vertical maps to be isomorphisms is that the Goodwillie-Weiss spectral sequence collapses along the anti-diagonal (Theorem 6.1). Theorem A is then a consequence of such collapse results. To put it differently, Theorem A says that the kernel of the vertical map  $\pi_0(K)/\sim_n \rightarrow \pi_0(T_{n+1}(K))$  can only have  $p$ -power torsion for  $p < n - 1$ . This result is modest in comparison to the conjecture that the two towers are isomorphic over  $\mathbb{Z}$ . Nevertheless, it seems to be the first non-rational computation about the Goodwillie-Weiss tower beyond low degrees ( $n \leq 3$  [9]).

We also note that, after tensoring with  $\mathbb{Q}$ , our result gives a construction of an isomorphism

$$(\pi_0(K)/\sim_n) \otimes \mathbb{Q} \cong \bigoplus_{s \leq n} \mathcal{A}_s^I \otimes \mathbb{Q}$$

that is purely homotopical and does not rely on transcendental techniques like Kontsevich's integral. However, our splitting is not explicit, whereas Kontsevich's integral gives a preferred choice of splitting over the complex numbers.

Theorem A is a corollary of more general results about the Goodwillie-Weiss tower for knots in  $\mathbb{R}^d$  – namely Theorem B below – as we now explain. This tower converges to the space of knots when  $d \geq 4$ , in the sense that the map from  $\text{emb}_c(\mathbb{R}, \mathbb{R}^d)$  to the homotopy limit of the tower is a weak equivalence. For  $d = 3$ , the case of classical knots, the tower is known not to converge to the knot space. (For example,  $\pi_0 \text{emb}_c(\mathbb{R}, \mathbb{R}^3)$  is countable whereas  $\pi_0 T_\infty \text{emb}_c(\mathbb{R}, \mathbb{R}^3)$  is uncountable.)

The spectral sequence associated to the Goodwillie-Weiss tower has  $E^1$ -page consisting of the homotopy groups of the layers of the tower, just like in the case  $d = 3$ . *Rationally*, this spectral sequence is well understood: from the work of Arone, Lambrechts, Volić and Turchin [1] we know that there is rational collapse at the  $E^2$ -page when  $d \geq 4$ . Our main technical result gives a vanishing range of differentials, at a prime  $p$ :

**Theorem B.** *Let  $p$  be a prime number. The differential*

$$d_{s,t}^r : E_{s,t}^r \rightarrow E_{s-r,t+r-1}^r$$

*vanishes  $p$ -locally if  $r - 1$  is not a multiple of  $(p - 1)(d - 2)$  and if*

$$t < 2p - 2 + (s - 1)(d - 2) .$$

Together with a result about triviality of extensions, this gives a computation of some homotopy groups of  $T_n \text{emb}_c(\mathbb{R}, \mathbb{R}^d)$ . The range improves as the prime or the codimension increases.

**Theorem C.** *Let  $n$  be a positive integer. For any prime number  $p$  satisfying  $n \leq (p - 1)(d - 2) + 3$ , there is a (non-canonical) isomorphism*

$$\pi_i T_n \text{emb}_c(\mathbb{R}^1, \mathbb{R}^d) \otimes \mathbb{Z}_{(p)} \cong \bigoplus_{t-s=i} E_{-s,t}^2(T_n) \otimes \mathbb{Z}_{(p)}$$

*for  $i \leq 2p - 6 + 2(d - 2)$ .*

The proof of Theorem B stands on the relation between the Goodwillie-Weiss tower and the little disks operad [25, 26, 13, 4], and the existence of Galois symmetries on the little disks operad that we established in [3]. Our main message in this paper is that by combining these two observations one obtains an interesting Galois action on the Goodwillie-Weiss tower. The numerology in Theorem B may seem esoteric at first, but it is a consequence of the action: the inequality has something

to do with the range over which the Galois action factors through the cyclotomic character.

We also prove a homological statement in Section 7 which, perhaps unsurprisingly, is stronger. It is also easier to prove, even though the strategy is the same. In [25], Sinha constructs a cosimplicial space  $K^\bullet$  which is a model for the Goodwillie-Weiss tower. Associated to the said cosimplicial space, and given a ring  $R$  of coefficients, there is a homological Bousfield-Kan spectral sequence  $E^*(R)$ . Modulo convergence issues, this spectral sequence is related to  $H_*(\overline{\text{emb}}_c(\mathbb{R}^1, \mathbb{R}^d), R)$  where  $\overline{\text{emb}}_c(\mathbb{R}^1, \mathbb{R}^d)$  denotes the homotopy fiber of the map

$$\text{emb}_c(\mathbb{R}^1, \mathbb{R}^d) \rightarrow \text{imm}_c(\mathbb{R}^1, \mathbb{R}^d).$$

Our main theorem about the homology spectral sequence is the following.

**Theorem D.** *Let  $p$  be a prime number. The only possibly non-trivial differentials in the spectral sequence  $E^*(\mathbb{Z}_p)$  are  $d_{1+n(d-1)(p-1)}$  for  $n \geq 0$ .*

This theorem was announced in [18]. Asymptotically, we also recover the collapse results for  $R = \mathbb{Q}$ , first established by Lambrechts-Turchin-Volić in [22] using the formality of the little disks operad.

**Acknowledgments.** We wish to thank Peter Teichner, Yuqing Shi and Danica Kosanović for valuable discussions.

## 1. BOUSFIELD LOCALIZATION

In this section, we recall some facts about localization of spaces. We denote the  $p$ -adic integers by  $\mathbb{Z}_p$ . Let  $E$  be a homology theory on spaces.

**Definition 1.1.** A space  $X$  is  $E$ -local if for any map  $f : U \rightarrow V$  such that  $E(f)$  is an isomorphism, the map induced by precomposition

$$\text{Map}(V, X) \rightarrow \text{Map}(U, X)$$

is an isomorphism.

**Theorem 1.2** (Bousfield [6]). *There exists an endofunctor  $L_E$  of the category of spaces equipped with a natural transformation  $\text{id} \rightarrow L_E$  satisfying the following two conditions.*

- (1) *For each  $X$  the space  $L_E(X)$  is  $E$ -local.*
- (2) *The map  $X \rightarrow L_E(X)$  induces an isomorphism in  $E$ -homology.*

*Moreover, the functor  $L_E$  is uniquely defined up to weak equivalence by these two properties.*

In the case where  $E$  is the homology theory  $H_*(-, \mathbb{F}_p)$ , the functor  $L_E$  will be denoted  $L_p$  and called  $p$ -completion, following common practice. This is justified by the next proposition.

**Proposition 1.3.** *If  $X$  is a simply connected space whose homotopy groups are finitely generated, the map  $X \rightarrow L_p X$  induces the canonical map  $\pi_*(X) \rightarrow \pi_*(X) \otimes \mathbb{Z}_p$  on homotopy groups.*

*Proof.* See [6, Proposition 4.3 (ii)]. □

We will need two facts about this construction.

**Proposition 1.4.** *The  $p$ -completion functor  $L_p$  commutes with products up to homotopy. More precisely, the canonical map*

$$L_p(X \times Y) \rightarrow L_p(X) \times L_p(Y)$$

*is an equivalence.*

*Proof.* The canonical map  $X \times Y \rightarrow L_p(X) \times L_p(Y)$  is a mod  $p$  homology equivalence by the Künneth isomorphism. Moreover, the target is easily seen to be  $p$ -complete. Therefore, by uniqueness of the  $p$ -completion,  $L_p(X) \times L_p(Y)$  must be the  $p$ -completion of  $X \times Y$ .  $\square$

The following proposition shows that under some mild assumptions on  $X$ , the homology of  $X$  with coefficients in the  $p$ -adic integers  $\mathbb{Z}_p$  can be recovered from the  $p$ -completion. We denote by  $C_*$  the standard singular chains functor from spaces to chain complexes.

**Proposition 1.5.** *Let  $X$  be a space whose homology with coefficients in  $\mathbb{Z}_p$  is degreewise finitely generated. Then the composite*

$$C_*(X, \mathbb{Z}_p) \rightarrow \lim_n C_*(X, \mathbb{Z}/p^n) \rightarrow \lim_n C_*(L_p X, \mathbb{Z}/p^n)$$

*is a quasi-isomorphism.*

*Proof.* We show that each of the two maps is a quasi-isomorphism. First, we observe that each of the two towers is a tower of epimorphisms. It follows that the limit agrees with the homotopy limit. In order to prove that the second map is a quasi-isomorphism, it is enough to prove that the map  $f : X \rightarrow L_p X$  is a mod  $p^n$  homology equivalence for all  $n$ . This follows by an easy induction using the Bockstein long exact sequence

$$\dots \rightarrow H_i(-, \mathbb{Z}/p^n) \rightarrow H_i(-, \mathbb{Z}/p^{n+1}) \rightarrow H_i(-, \mathbb{Z}/p) \rightarrow H_{i-1}(-, \mathbb{Z}/p^n) \rightarrow \dots$$

We now show that the map

$$C_*(X, \mathbb{Z}_p) \rightarrow \lim_n C_*(X, \mathbb{Z}/p^n)$$

is a quasi-isomorphism. The homology groups of the source are, by definition, the homology groups of  $X$  with  $\mathbb{Z}_p$  coefficients and the homology groups of the target are

$$H_i(\lim_n C_*(X, \mathbb{Z}/p^n)) \cong \lim_n H_i(X, \mathbb{Z}/p^n).$$

This isomorphism follows from a Milnor short exact sequence argument (the assumption made on  $X$  implies that the mod  $p^n$  homology is degreewise finite and thus the  $\lim^1$ -term vanishes by the Mittag-Leffler criterion). Thus we have to prove that the canonical map

$$H_i(X, \mathbb{Z}_p) \rightarrow \lim_n H_i(X, \mathbb{Z}/p^n)$$

is an isomorphism. This map can be factored as

$$H_i(X, \mathbb{Z}_p) \rightarrow \lim_n H_i(X, \mathbb{Z}_p) \otimes \mathbb{Z}/p^n \rightarrow \lim_n H_i(X, \mathbb{Z}/p^n).$$

The first of these two maps is an isomorphism under the finite type assumption. The second map is also an isomorphism. This can be seen as follows: using the universal coefficients theorem and the left exactness of the limit functor, we have an exact sequence

$$0 \rightarrow \lim_n H_i(X, \mathbb{Z}_p) \otimes \mathbb{Z}/p^n \rightarrow \lim_n H_i(X, \mathbb{Z}/p^n) \rightarrow \lim_n \operatorname{Tor}^{\mathbb{Z}_p}(\mathbb{Z}/p^n, H_{i-1}(X, \mathbb{Z}_p)).$$

But, for any finitely generated  $\mathbb{Z}_p$ -module  $A$ , the group  $\lim_n \mathrm{Tor}(\mathbb{Z}/p^n, A)$  is zero.  $\square$

**Remark 1.6.** This proposition may be false if the homology of  $X$  is not finitely generated. As a counter-example, one can take  $X$  to be the rationalization of  $S^1$ . On the one hand, its  $p$ -completion is contractible. On the other hand,  $H_1(X, \mathbb{Z}_p)$  is isomorphic to  $\mathbb{Q}_p$ .

## 2. A GROTHENDIECK-TEICHMÜLLER ACTION ON THE LITTLE DISKS OPERADS

The Grothendieck-Teichmüller group  $GT_p$  introduced in [12] is a profinite group equipped with a surjective map

$$\chi : GT_p \rightarrow \mathbb{Z}_p^\times$$

called the cyclotomic character. Given a  $\mathbb{Z}_p$ -module  $M$ , we can give it a  $GT_p$ -action using the formula

$$\gamma.m := \chi(\gamma)^n m .$$

We call this action the weight  $n$  cyclotomic action.

**Proposition 2.1.** *Let  $M$  be a  $\mathbb{Z}_p$ -module equipped with a  $GT_p$ -action. Suppose that  $M$  fits in a short exact sequence of  $GT_p$ -modules*

$$0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0 .$$

*If the  $GT_p$ -action on  $M$  is cyclotomic of weight  $n$ , then that is also the case for the  $GT_p$ -action on  $M'$  and  $M''$ .*

*Proof.* This is an immediate verification.  $\square$

**Proposition 2.2.** *Let  $M$  and  $N$  be two  $\mathbb{Z}_p$ -modules equipped, respectively, with the weight  $m$  and weight  $n$  cyclotomic  $GT_p$ -action. If  $m - n$  is not a multiple of  $p - 1$ , any  $GT_p$ -equivariant map*

$$f : M \rightarrow N$$

*must be the zero map.*

*Proof.* Let us assume that  $m - n$  is not a multiple of  $p - 1$ . Let  $x$  be any element of  $M$ . If  $f$  is a  $GT_p$ -equivariant map, we must have

$$\chi(\gamma)^m f(x) = \chi(\gamma)^n f(x)$$

for every  $\gamma$ . In particular, since  $\chi$  is surjective, for any unit  $u$  of  $\mathbb{Z}_p$  we have that  $(u^{n-m} - 1) \cdot f(x) = 0$ . Hence, it is enough to find  $u$  such that  $u^{n-m} - 1$  is also a unit. Take a generator  $v$  of  $\mathbb{F}_p^\times$ . Then  $v^{n-m} \neq 1$  if  $n - m$  is not a multiple of  $p - 1$  (since  $\mathbb{F}_p^\times$  is cyclic of order  $p - 1$ ). Let  $u$  be a lift of  $v$ . Then  $u^{n-m} - 1$  is congruent to  $v^{n-m} - 1$  modulo  $p$  and so it is non-zero modulo  $p$ , i.e. it is a unit.  $\square$

The next theorem plays a key role in what follows; it equips the  $p$ -completion of the little disks operad with an action of  $GT_p$ . We denote by  $E_d$  the little disks operad of dimension  $d$ . It will be convenient to view it as an  $\infty$ -operad (a dendroidal space satisfying the Segal condition). Applying  $L_p$  objectwise, we obtain a dendroidal space  $L_p E_d$  which, by Proposition 1.4, also satisfies the Segal condition. For more details, see [3].

**Theorem 2.3.** *Let  $d$  be an integer with  $d \geq 3$ . There exists a  $GT_p$ -action on  $L_p E_d$  such that the induced action on  $H_{d-1}(L_p E_d(2), \mathbb{Z}_p)$  is the cyclotomic action of weight 1. Moreover, if we give  $E_1$  the trivial  $GT_p$ -action, the canonical map  $E_1 \rightarrow L_p E_d$  is  $GT_p$ -equivariant.*

*Proof.* This is done in [3]. We recall briefly how this construction goes. For a space  $X$ , we denote by  $X^\wedge$  its pro- $p$ -completion. This is a pro-object in  $p$ -finite spaces (a space is  $p$ -finite if it has finitely many components and if each component is truncated with finite  $p$ -groups as homotopy groups) that receives a map from  $X$  and is initial for this property. There is a close relationship between this functor and the  $p$ -completion functor  $X \mapsto L_p(X)$ : when  $X$  is a nilpotent space of finite  $\mathbb{F}_p$ -type (i.e. all its  $\mathbb{F}_p$ -homology groups are finite dimensional) the  $p$ -completion  $L_p(X)$  coincides with the homotopy limit of the pro-object  $X^\wedge$ . This homotopy limit is denoted  $\mathbb{R}Mat$  in [3] and is the homotopy right adjoint of the pro- $p$ -completion functor.

By work of Drinfel'd, there exists a non-trivial action of  $GT_p$  on  $E_2^\wedge$ . Using a version of the Dunn-Lurie additivity theorem established in [5] one can produce an action of  $GT_p$  on  $E_d^\wedge$  by writing it as a certain tensor product of  $(E_2)_{\mathbb{F}_p}^\wedge$  with the Drinfel'd action and  $E_{d-2}^\wedge$  with the trivial action. Moreover, with respect to this action, the map  $E_1^\wedge \rightarrow E_d^\wedge$  is  $GT_p$ -equivariant. Applying the functor  $\mathbb{R}Mat$ , we obtain the asserted  $GT_p$ -equivariant map  $E_1 \simeq L_p E_1 \rightarrow L_p E_d$  using the fact that, for  $d \geq 3$ , the spaces  $E_d(k)$  are nilpotent and of finite type.

The statement that the action is cyclotomic of weight 1 on  $H_{d-1}(L_p E_d(2), \mathbb{Z}_p)$  follows from [3, Proof of Theorem 7.1] where it is shown that the composite

$$GT_p \rightarrow \text{Aut}^h(L_p E_d) \rightarrow \text{Aut}^h(L_p E_d(2)) \cong \mathbb{Z}_p^\times$$

coincides with the cyclotomic character. □

From this, we can fully understand the induced  $GT_p$ -action on homology.

**Proposition 2.4.** *For any  $k$ , the  $GT_p$ -action on  $H_n(d-1)(E_d(k), \mathbb{Z}_p)$  is cyclotomic of weight  $n$ .*

*Proof.* The homology of  $E_d$  is the operad of  $d$ -Poisson algebras; it is generated by a commutative product of degree 0, a Lie bracket of degree  $(d-1)$ , the latter being a derivation with respect to the former in both variables. Since this operad is generated by operations of arity 2, it follows that the homology  $H_*(E_d(k), \mathbb{Z}_p)$  is a quotient of tensor powers of  $H_*(E_d(2), \mathbb{Z}_p)$ . The statement of the proposition holds for  $k=2$  by Theorem 2.3 and the general case follows from Proposition 2.1. □

### 3. THE GOODWILLIE-WEISS TOWER AND $p$ -COMPLETION

**3.1. Generalities about  $p$ -completion of towers.** For a tower of based spaces and basepoint preserving maps,

$$\cdots \rightarrow T_n \rightarrow T_{n-1} \rightarrow \cdots \rightarrow T_0$$

there is an associated spectral sequence [7, IX.4] whose  $E^1$  page is

$$E_{-s,t}^1(T) = \pi_{t-s} \text{hofiber}(T_s \rightarrow T_{s-1})$$

and the first differential  $d^1 : E_{-s,t}^1(T) \rightarrow E_{-(s+1),t}^1(T)$  is the composite

$$\pi_{t-s} \text{hofiber}(T_s \rightarrow T_{s-1}) \rightarrow \pi_{t-s} T_s \rightarrow \pi_{t-s-1} \text{hofiber}(T_s \rightarrow T_{s-1})$$

where the second map is the boundary homomorphism.

We assume that each  $T_i$  has abelian homotopy groups (including  $\pi_0$ ) and that the induced maps on homotopy groups are group homomorphisms (e.g. a tower of 2-fold loop spaces). This is the case in our examples below, and having this hypothesis avoids subtleties which arise when defining this spectral sequence in general, when  $\pi_0$  and  $\pi_1$  are not abelian groups. The differential  $d^r$  has degree  $(-r, r-1)$ .

Let us briefly discuss convergence. For details, see [7, IX]. Let  $F^s \pi_*(\text{holim} T)$  be the image of the map  $\pi_*(\text{holim} T) \rightarrow \pi_* T_s$ . Using the Milnor short exact sequence, we see that, up to a  $\lim^1$ -term, the group  $\pi_*(\text{holim} T)$  is the limit of the tower of epimorphisms

$$(3.1) \quad \cdots \rightarrow F^s \pi_*(\text{holim} T) \rightarrow F^{s-1} \pi_*(\text{holim} T) \rightarrow \cdots \rightarrow F^0 \pi_*(\text{holim} T)$$

(indeed the limit of this tower can easily be identified with  $\lim_s \pi_* T_s$ ) and there are inclusions

$$\ker[F^s \pi_*(\text{holim} T) \rightarrow F^{s-1} \pi_*(\text{holim} T)] \subset E_{-s, s+*}^\infty(T).$$

For these inclusions to be isomorphisms, a sufficient condition is that

$$\lim_r^1 E_{-s, t}^r$$

vanishes [7, IX.5.4]. Under this hypothesis, the term  $\lim_s^1 \pi_* T_s$  also vanishes, and so the spectral sequence converges in the sense that the limit of the tower (3.1) is  $\pi_*(\text{holim} T)$  and the associated graded of this tower is given by the  $E^\infty$ -page. The vanishing of  $\lim_r^1 E_{-s, t}^r$  happens, for example, if the connectivity of the layers of the tower  $T$  goes to  $\infty$  as  $n \rightarrow \infty$ .

Before we get to our main example, we need to discuss the effect of applying  $p$ -completion  $L_p$  to a certain tower of spaces. Suppose  $\{X_n\}_{n \in \mathbb{N}}$  is a tower of simply connected based spaces with finitely generated homotopy groups. Assume further that the layers are also simply connected and consider the corresponding tower  $T = \{\Omega^2 X_n\}$  of two-fold loop spaces and maps. We wish to compare the tower  $T$  with a tower  $T(\mathbb{Z}_p)$

$$\cdots \rightarrow \Omega^2 L_p X_n \rightarrow \Omega^2 L_p X_{n-1} \rightarrow \cdots \rightarrow \Omega^2 L_p X_0$$

and an analogous spectral sequence whose  $E^1$ -page is

$$E_{-s, t}^1(T(\mathbb{Z}_p)) = \pi_{t-s} \Omega^2 \text{hofiber}(L_p X_s \rightarrow L_p X_{s-1}).$$

A warning must be issued at this point: taking homotopy fibers does not commute with  $L_p$  in general. However, under the simple connectivity assumption, we have that

$$\text{hofiber}(L_p X_s \rightarrow L_p X_{s-1}) \simeq L_p \text{hofiber}(X_s \rightarrow X_{s-1}).$$

Therefore,

**Proposition 3.1.** *For each positive integer  $r$ , the map*

$$E_{-s, t}^r(T) \rightarrow E_{-s, t}^r(T(\mathbb{Z}_p))$$

*is given by algebraic  $p$ -completion, i.e., tensoring with  $\mathbb{Z}_p$ .*



*Proof.* For  $r = 1$ , this follows from what was just said and the fact that if a space  $Y$  has homotopy groups that are finitely generated abelian, then the map

$$\pi_* Y \rightarrow \pi_* L_p Y$$

is a  $p$ -completion, in the sense that it is isomorphic to the canonical map

$$\pi_* Y \rightarrow \pi_* Y \otimes \mathbb{Z}_p .$$

For a finite  $r$ , the result follows by the exactness the functor  $A \mapsto A \otimes \mathbb{Z}_p$ .  $\square$

Now, suppose that we are in a situation for which there is convergence of the spectral sequence of the tower  $T$ , e.g. if the connectivity of the layers of  $T$  grows to  $\infty$ . Then we have a tower approximating  $\pi_*(\text{holim}T(\mathbb{Z}_p))$

$$(3.2) \quad \cdots \rightarrow F_p^s \rightarrow F_p^{s-1} \rightarrow \cdots \rightarrow F_p^0$$

where  $F_p^s$  is the image of  $\pi_*(\text{holim}T(\mathbb{Z}_p)) \rightarrow \pi_* T(\mathbb{Z}_p)_s$ , and satisfying

$$E_{-s, s+*}^\infty(T(\mathbb{Z}_p)) \cong \ker(F_p^s \rightarrow F_p^{s-1}) .$$

This is related to the tower  $\{F^s\}$  which approximates  $\pi_*(\text{holim}T)$ . Namely, the tower (3.2) is a  $p$ -completion of the tower (3.1).

**Remark 3.2.** Without convergence hypothesis, the tower (3.2) may *not* be a  $p$ -completion of the tower (3.1). This is because  $p$ -completion can fail to commute with infinite limits. For the same reason, the map

$$\pi_*(\text{holim}T) \rightarrow \pi_*(\text{holim}T(\mathbb{Z}_p))$$

may not be a  $p$ -completion.

**3.2. The Goodwillie-Weiss tower for long knots.** Let  $d \geq 3$ . The Goodwillie-Weiss tower approximating the space of long knots  $\text{emb}_c(\mathbb{R}^1, \mathbb{R}^d)$

$$\cdots \rightarrow T_k \text{emb}_c(\mathbb{R}^1, \mathbb{R}^d) \rightarrow \cdots \rightarrow T_2 \text{emb}_c(\mathbb{R}^1, \mathbb{R}^d)$$

is a tower of 2-fold loop spaces. We briefly explain this here, following [4]. The  $k$ -th term,  $k \geq 2$ , can be described via a homotopy fiber sequence

$$T_k \text{emb}_c(\mathbb{R}^1, \mathbb{R}^d) \rightarrow \Omega \text{InjLin}(\mathbb{R}^1, \mathbb{R}^d) \xrightarrow{\Omega f} \Omega \text{Map}^h(E_1, E_d)_{\leq k}$$

where  $\text{InjLin}$  denotes the space of injective linear maps, and  $\text{Map}^h(E_1, E_d)_{\leq k}$  denotes the space of derived maps between  $k$ -truncated operads. (That is, we restrict the operads to operations with at most  $k$  inputs.) The map  $f$  is essentially an evaluation map (see [4] for a precise description using configuration categories).

The space  $\text{InjLin}(\mathbb{R}^1, \mathbb{R}^d)$  is weakly equivalent to  $\text{Map}^h(E_1, E_d)_{\leq 2}$  and with that in mind it is not hard to see that  $f$  is a section of the truncation map

$$g_k : \text{Map}^h(E_1, E_d)_{\leq k} \rightarrow \text{Map}^h(E_1, E_d)_{\leq 2} .$$

Therefore, we have a weak equivalence

$$T_k \text{emb}_c(\mathbb{R}^1, \mathbb{R}^d) \simeq \Omega^2 \text{hofiber}(g_k) .$$

Moreover, the map

$$T_{k+1} \text{emb}_c(\mathbb{R}^1, \mathbb{R}^d) \rightarrow T_k \text{emb}_c(\mathbb{R}^1, \mathbb{R}^d)$$

can be identified with the double loop map of the obvious map

$$\text{hofiber}(g_{k+1}) \rightarrow \text{hofiber}(g_k) .$$

In order to simplify notation, we denote the tower  $k \mapsto \Omega^2 \text{hofiber}(g_k)$  by  $T$ .

For a positive integer  $n$ , we write  $\underline{n}$  for the finite set  $\{1, \dots, n\}$ . The first page of the spectral sequence associated to  $T$  has the following description: whenever  $s > 2$ , we have

$$E_{-s,t}^1(T) = \pi_t \text{thofiber}(S \subset \underline{s} \mapsto \text{emb}(S, \mathbb{R}^d))$$

where thofiber means the total homotopy fiber of the cube; when  $s \leq 2$ , we have  $E_{-s,t}^1(T) = 0$ . This identification follows from:

**Proposition 3.3** (Göppel [16]). *Let  $P$  be an operad having weakly contractible  $P(0)$  and  $P(1)$ . Let us fix a map  $E_1 \rightarrow P$  that we use as a base point for each of the spaces  $\text{Map}^h(E_1, P)_{\leq n}$ . For  $n \geq 2$ , there is a homotopy fiber sequence*

$$\Omega^{n-2} \text{thofiber}(S \subset \underline{n} \mapsto P(S)) \rightarrow \text{Map}^h(E_1, P)_{\leq n} \rightarrow \text{Map}^h(E_1, P)_{\leq n-1}$$

where the  $n$ -cube  $S \mapsto P(S)$  is given by compositions with 0-arity operations.

The homotopy groups of this total homotopy fiber are well-understood. We record this in the following two elementary and well-known statements.

**Lemma 3.4.** *Let  $X$  and  $Y$  be two  $(n-1)$  dimensional cubes. Let  $p : X \rightarrow Y$  and  $s : Y \rightarrow X$  be two maps of cubes with  $s$  a homotopy section of  $p$  (i.e.  $p \circ s$  is homotopic to the identity). Then, for any  $k \geq 1$ , if we denote by  $C$  the  $n$ -dimensional cube given by the map  $p$ , we have*

$$\pi_k(\text{thofiber}(C)) \cong \ker(\pi_k(\text{thofiber}(X)) \rightarrow \pi_k(\text{thofiber}(Y)))$$

where the map on the right hand side is the map induced by  $p$ .

*Proof.* In general, there is a fiber sequence

$$\text{thofiber}(C) \rightarrow \text{thofiber}(X) \xrightarrow{p} \text{thofiber}(Y)$$

and the existence of the map  $s$  implies that the long exact sequence of homotopy groups splits into short exact sequences:

$$0 \rightarrow \pi_k(\text{thofiber}(C)) \rightarrow \pi_k(\text{thofiber}(X)) \rightarrow \pi_k(\text{thofiber}(Y)) \rightarrow 0$$

as claimed. □

**Proposition 3.5.** *We have*

$$\pi_* \text{thofiber}(S \subset \underline{n} \mapsto P(S)) \cong \bigcap_{i=0}^{n-1} (\ker \pi_*(s^i) : \pi_* P(n) \rightarrow \pi_* P(n-1))$$

where the maps  $s^i : P(n) \rightarrow P(n-1)$  are the  $n$  edges of the cube originating from  $P(n)$ .

*Proof.* Let us call  $C$  the cube under consideration. Let  $z$  be any point of the space  $P(0)$  (the choice does not matter since this space is contractible). The map  $s^i$  is then defined as operadic composition with  $z$  in the  $(i+1)$ -st input. The cube  $C$  can be viewed as a map of  $(n-1)$ -cubes  $X \rightarrow Y$  with  $X(S) = P(S \cup \{n\})$  and  $Y(S) = P(S)$  for  $S \subset \underline{n-1}$ . The map from  $X$  to  $Y$  is induced by operadic composition with  $z$  in the input labelled  $n$ .

Recall that we have chosen a map  $E_1 \rightarrow P$ . Let  $x$  in  $P(2)$  be the image of any point in  $E_1(2)$  (again the choice is irrelevant). Using operadic composition with  $x$  we obtain a homotopy section  $Y \rightarrow X$  as in the previous lemma. By the previous lemma, we deduce that  $\pi_* \text{thofiber}(C)$  is the kernel of the map induced by  $s^{n-1}$  on

$\pi_* \text{thofiber}(X)$ . Iterating this reasoning  $(n - 1)$  more times, we obtain the desired result.  $\square$

We now fix a prime  $p$ . As in the discussion in the beginning of the section, we want to investigate the tower

$$\cdots \rightarrow \Omega^2 L_p \text{hofiber}(g_k) \rightarrow \Omega^2 L_p \text{hofiber}(g_{k-1}) \rightarrow \cdots$$

In line with the notation of the previous subsection, we denote this tower by  $T(\mathbb{Z}_p)$ . We also introduce another tower – denoted  $T'(\mathbb{Z}_p)$  – of the form

$$\cdots \rightarrow \Omega^2 \text{hofiber}(g'_k) \rightarrow \Omega^2 \text{hofiber}(g'_{k-1}) \rightarrow \cdots$$

where  $g'_k$  is the truncation map for the operad  $L_p E_d$  :

$$g'_k : \text{Map}^h(E_1, L_p E_d)_{\leq k} \rightarrow \text{Map}^h(E_1, L_p E_d)_{\leq 2} .$$

**Lemma 3.6.** *The canonical map  $T(\mathbb{Z}_p) \rightarrow T'(\mathbb{Z}_p)$  is a weak equivalence of towers. As such, it induces an isomorphism of the associated spectral sequences.*

*Proof.* We compare the layers. According to Propositions 3.3 and 3.5, the  $n^{\text{th}}$  layer of  $T'(\mathbb{Z}_p)$  has homotopy groups

$$\bigcap_{i=0}^{n-1} (\ker \pi_*(s^i) : (\pi_* E_d(n)) \otimes \mathbb{Z}_p \rightarrow (\pi_* E_d(n-1)) \otimes \mathbb{Z}_p) .$$

By Proposition 3.1, the homotopy groups of the layers of  $T(\mathbb{Z}_p)$  are the  $p$ -completion of the homotopy groups of the layers of  $T$ . The homotopy groups of the layers of  $T$  are also identified using Propositions 3.3 and 3.5. Taking  $p$ -completion amounts to tensoring the homotopy groups with  $\mathbb{Z}_p$ , which is an exact functor since  $\mathbb{Z}_p$  is torsion-free, and so the result follows.  $\square$

The important corollary of the above discussion is the following.

**Corollary 3.7.** *The tower  $T'(\mathbb{Z}_p)$  has an action of  $GT_p$ . Therefore, the spectral sequence  $E^r_{-s,t}(T(\mathbb{Z}_p)) \cong E^r_{-s,t}(T'(\mathbb{Z}_p))$  has an action of  $GT_p$ .*

*Proof.* This is a straightforward consequence of Theorem 2.3.  $\square$

**Remark 3.8.** In [19, Appendix D], Kassel and Turaev construct a Galois action on the Vassiliev tower over  $\mathbb{Q}_p$ . As we show later, the Vassiliev tower over  $\mathbb{Q}_p$  is isomorphic to the Goodwillie-Weiss tower  $\pi_0 T(\mathbb{Q}_p)$  (defined in the next section). But we do not know whether the action from Corollary 3.7 agrees with the one of Kassel and Turaev.

It is also not absurd to wonder whether there is a relation between the tower  $T'(\mathbb{Z}_p)$  and the *profinite knots* of Furusho [14], who also come with an action of the Grothendieck-Teichmüller group.

**Remark 3.9.** There is a closely related tower

$$\cdots \rightarrow \bar{T}_k \rightarrow \cdots \rightarrow \bar{T}_2$$

approximating the space

$$\overline{\text{emb}}_c(\mathbb{R}^1, \mathbb{R}^n) = \text{hofiber}(\text{emb}_c(\mathbb{R}^1, \mathbb{R}^n) \rightarrow \Omega \text{InjLin}(\mathbb{R}^1, \mathbb{R}^d))$$

and whose  $k$ -term is

$$\bar{T}_k = \text{hofiber}(T_k \rightarrow \Omega \text{InjLin}(\mathbb{R}^1, \mathbb{R}^d)) \simeq \Omega^2 \text{Map}^h(E_1, E_d)_{\leq k} .$$

This tower is identified (Sinha [25]) with the tower associated to a pointed cosimplicial space  $K^\bullet$

$$\cdots \rightarrow \operatorname{holim}_{\Delta_{\leq k}} K^\bullet \rightarrow \operatorname{holim}_{\Delta_{\leq k-1}} K^\bullet \rightarrow \cdots \rightarrow \operatorname{holim}_{\Delta_{\leq 0}} K^\bullet = K^0$$

The first page of the spectral sequence associated to this tower  $\bar{T}$  agrees with that of the tower  $T$ , except on the column  $s = 2$ . All the discussion above applies to the tower  $\bar{T}$  without any additional difficulty. In section 7 we study the Bousfield-Kan homology spectral sequence of this cosimplicial space.

#### 4. MAIN THEOREM

In this section, we prove the following Theorem from which Theorem B of the introduction is an easy consequence.

**Theorem 4.1.** *Let  $p$  be a prime number. The differential*

$$d_{-s,t}^r : E_{-s,t}^r(T(\mathbb{Z}_p)) \rightarrow E_{-s-r,t+r-1}^r(T(\mathbb{Z}_p))$$

*vanishes if  $r - 1$  is not a multiple of  $(p - 1)(d - 2)$  and if*

$$t < 2p - 2 + (s - 1)(d - 2) .$$

We begin by recalling the theorem of Hilton-Milnor in a form that we will need. Given a finite set  $R$ , we denote by  $W(R)$  a set of Lie words in the finite set  $R$  that forms a basis for the free Lie algebra on  $R$ . For  $w \in W(R)$  we write  $|w|$  for its length.

**Theorem 4.2** (Hilton-Milnor). *There is a weak equivalence*

$$\prod'_{w \in W(R)} \Omega S^{|w|(d-2)+1} \xrightarrow{\simeq} \Omega \left( \bigvee_R S^{d-1} \right)$$

*where the symbol  $\prod'$  stands for the weak product (union of finite products).*

Let  $T(\mathbb{Q}_p)$  be the tower analogous to  $T(\mathbb{Z}_p)$  obtained by replacing all instances of  $L_p Z$ ,  $Z$  a space, by their rationalizations  $(L_p Z)_{\mathbb{Q}}$ . As in the previous section, we have a  $GT_p$ -action on  $T(\mathbb{Q}_p)$  and so a  $GT_p$ -action on the associated spectral sequence  $E(T(\mathbb{Q}_p))$  (c.f. Corollary 3.7). Note also that by exactness of the functor  $- \otimes_{\mathbb{Z}_p} \mathbb{Q}_p$ , we have the following isomorphisms for any  $r, s$  and  $t$

$$E_{-s,t}^r(T(\mathbb{Q}_p)) \cong E_{-s,t}^r(T(\mathbb{Z}_p)) \otimes \mathbb{Q} \cong E_{-s,t}^r(T) \otimes \mathbb{Q}_p$$

the first of which is  $GT_p$ -equivariant.

**Theorem 4.3.** *Let  $d \geq 3$ . The  $\mathbb{Q}_p$ -vector space*

$$E_{-s,t}^1(T(\mathbb{Q}_p))$$

*is zero unless  $t = n(d - 2) + 1$  for some  $n \geq s - 1$ . If  $t = n(d - 2) + 1$ , the  $GT_p$ -action is cyclotomic of weight  $n$ .*

*Proof.* We assume  $s > 2$ , otherwise  $E_{-s,t}^1(T(\mathbb{Q}_p))$  is zero. Then

$$E_{-s,t}^1(T(\mathbb{Q}_p)) = \pi_t \text{thofiber} (S \subset \underline{s} \mapsto L_p \operatorname{emb}(S, \mathbb{R}^d)) \otimes \mathbb{Q} .$$

Regard the  $s$ -cube  $S \mapsto L_p \operatorname{emb}(S, \mathbb{R}^d)$  as an  $(s - 1)$ -cube of maps

$$(R \subset \underline{s-1}) \mapsto \chi_R$$

where  $\chi_R$  is the map  $L_p \text{emb}(R \cup \{s\}, \mathbb{R}^d) \rightarrow L_p \text{emb}(R, \mathbb{R}^d)$  forgetting the point labelled  $s$ . Then the total homotopy fiber of the original cube is identified with the total homotopy fiber of the  $(s-1)$ -cube  $R \mapsto \text{hofiber} \chi_R$ . This  $(s-1)$ -cube is identified with a cube

$$R \mapsto \phi_R := L_p(\vee_R S^{d-1}).$$

where the maps collapse wedge summands. This follows from the Fadell-Neuwirth fiber sequence

$$\vee_R S^{d-1} \rightarrow \text{emb}(R \cup \{s\}, \mathbb{R}^d) \rightarrow \text{emb}(R, \mathbb{R}^d)$$

and the fact that  $L_p$  commutes with taking homotopy pullbacks of simply connected spaces. By Proposition 3.5, the canonical projection

$$(4.1) \quad \text{thofiber}(\phi) \rightarrow L_p(\vee_{s-1} S^{d-1})$$

is injective on homotopy groups. Now, the  $GT_p$  action on  $E_{-s,t}^1(\mathbb{Q}_p)$  comes from a (basepoint preserving) action on the cube  $S \mapsto L_p \text{emb}(S, \mathbb{R}^d)$ , or equivalently, from an action on the cube  $\phi$ . As such, the projection map (4.1) is  $GT_p$ -equivariant. This means that to understand the action on  $E_{-s,t}^1(\mathbb{Q}_p)$  it is enough to understand the action on

$$\pi_t L_p(\vee_{s-1} S^{d-1}) \otimes \mathbb{Q} \cong \pi_t(\vee_{s-1} S^{d-1}) \otimes \mathbb{Q}_p.$$

Using Hilton-Milnor theorem (Theorem 4.2) or, alternatively, Milnor-Moore, the rational homotopy groups of a wedge of  $(d-1)$ -dim spheres are concentrated in degrees  $n(d-2)+1$ , for  $n \geq 1$ . The integer  $n$  records the *size* of the word in the Hilton-Milnor decomposition. In the total homotopy fiber of the  $(s-1)$ -cube  $R \mapsto \vee_R S^{d-1}$ ,  $n$  must be at least  $s-1$  since the generator  $\iota \in \pi_{d-1}(S^{d-1})$  of each wedge summand must occur at least once. This establishes the first part of the theorem. The second part is a consequence of the fact that the  $GT_p$ -action on  $\pi_{n(d-2)+1}(\vee S^{d-1}) \otimes \mathbb{Q}_p$  is cyclotomic of weight  $n$ . This is the content of Proposition 4.4 below when  $n=1$ . For higher  $n$ 's, these homotopy groups are generated by Whitehead products of elements in  $\pi_{d-1}$ , and since the  $GT_p$  action exists at the level of spaces it must be compatible with Whitehead products.  $\square$

**Proposition 4.4.** *The  $GT_p$  action on*

$$\pi_{d-1} L_p(\vee_k S^{d-1}) \cong \oplus_k \mathbb{Z}_p$$

*is cyclotomic of weight 1.*

*Proof.* For the duration of this proof, we will drop  $L_p$  from the notation and so, for a space  $X$ , we keep  $X$  as notation for  $L_p X$ .

By Proposition 2.4, the statement holds for  $k=1$ . For  $k > 1$ , set  $\ell = k+1$  and consider the commutative square

$$\begin{array}{ccc} \text{emb}(\underline{\ell}, \mathbb{R}^d) & \xrightarrow{\text{forget } k+1} & \text{emb}(\underline{k}, \mathbb{R}^d) \\ \downarrow & & \downarrow \\ \text{emb}(\underline{2}, \mathbb{R}^d) & \xrightarrow{\text{forget } 2} & \text{emb}(\underline{1}, \mathbb{R}^d) \end{array}$$

where the left map is induced by the inclusion  $f : \underline{2} \rightarrow \underline{\ell}$  given by  $f(1) = j$ ,  $f(2) = k+1$ ; and the right map is induced by the map  $\underline{1} \rightarrow \underline{k}$  selecting  $j$ . The induced map on horizontal homotopy fibers has the form

$$p_j : \vee_k S^{d-1} \rightarrow S^{d-1}$$

and it corresponds to collapsing all summands except the  $j$ -th one to the basepoint. All the (based) maps in the square are  $GT_p$ -equivariant since they correspond to composition maps in the operad  $E_d$  involving nullary operations, and so  $p_j$  is a  $GT_p$ -equivariant map.

Let  $\alpha$  be an element of  $GT_p$ . To describe the action of  $\alpha$  on the wedge  $\vee_k S^{d-1}$  we must describe, for each pair  $1 \leq i, j \leq k$ , the composition

$$S^{d-1} \xrightarrow{\text{incl}_i} \vee_k S^{d-1} \xrightarrow{\alpha} \vee_k S^{d-1} \xrightarrow{p_j} S^{d-1}$$

where  $\text{incl}_i$  means the inclusion of the  $i$ -th summand. Since  $p_j \circ \alpha = \alpha \circ p_j$ , this composite equals  $\alpha \circ p_j \circ \text{incl}_i$ , which is  $\alpha$  if  $i = j$  and is trivial otherwise.  $\square$

*Proof of Theorem 4.1.* Let  $W$  denote the total homotopy fiber of the  $(s-1)$ -cube  $R \mapsto \vee_R S^{d-1}$ , for  $s > 2$ . We look at the  $p$ -torsion in the homotopy groups of  $W$ . By the naturality in the Hilton-Milnor theorem (Theorem 4.2), it follows that

$$\Omega W \simeq \prod'_w \Omega S^{|w|(d-2)+1}$$

where  $w$  runs over the words in the letters  $1, \dots, s-1$  containing every letter *at least once*. As such, the smallest word in the product has length  $s-1$ , and the sphere of smallest dimension is  $S^{(s-1)(d-2)+1}$ .

By a famous result of Serre, the homotopy groups  $\pi_* S^\ell$  are  $p$ -torsion free for  $* \leq \ell + 2p - 4$  for  $\ell \geq 3$ . Therefore, the homotopy groups  $\pi_* W$  are  $p$ -torsion free whenever  $* \leq N$  where

$$N = (s-1)(d-2) + 2p - 3,$$

(remember  $d \geq 3$  and  $s \geq 3$  so that  $(s-1)(d-2) + 1 \geq 3$ ). And so, in the range  $t \leq N$ , we have an inclusion

$$E_{-s,t}^1(T(\mathbb{Z}_p)) \hookrightarrow E_{-s,t}^1(T(\mathbb{Z}_p)) \otimes \mathbb{Q} \cong E_{-s,t}^1(T(\mathbb{Q}_p)).$$

Therefore, by Theorem 4.3, in the range  $t \leq N$ , the group  $E_{-s,t}^1(T(\mathbb{Z}_p))$  is zero unless  $t = n(d-2) + 1$ , in which case  $GT_p$  acts with weight  $n$ . The same is true for the successive pages  $E_{-s,t}^r$  by Proposition 2.1. Proposition 2.2 completes the proof.  $\square$

We can now prove Theorem B of the introduction.

*Proof of Theorem B.* Recall that  $\mathbb{Z}_{(p)}$  denotes the ring of  $p$ -local integers. Proposition 3.1 gives an isomorphism of spectral sequences

$$E_{*,*}^*(T(\mathbb{Z}_p)) \cong E_{*,*}^*(T) \otimes_{\mathbb{Z}} \mathbb{Z}_p \cong (E_{*,*}^*(T) \otimes_{\mathbb{Z}} \mathbb{Z}_{(p)}) \otimes_{\mathbb{Z}_{(p)}} \mathbb{Z}_p$$

Then the result follows from the fact that a map  $f$  of  $\mathbb{Z}_{(p)}$ -modules vanishes if and only if  $f \otimes_{\mathbb{Z}_{(p)}} \mathbb{Z}_p$  vanishes.  $\square$

## 5. SOME CONSEQUENCES

**Corollary 5.1.** *The spectral sequence  $E_{-s,t}^*(T) \otimes_{\mathbb{Z}} \mathbb{Q}$  collapses at the  $E^2$ -page.*

*Proof.* Let  $d_{-s,t}^r$  be a differential in that spectral sequence with  $r > 2$ . Up to choosing  $p$  a large enough prime, we may assume that  $t < 2p - 2 + (s-1)(d-1)$  and that  $r-1$  is not a multiple of  $(p-1)(d-1)$ . Then this differential is zero in  $E_{-s,t}^r(T) \otimes_{\mathbb{Z}} \mathbb{Z}_{(p)}$  by Theorem B so it is also zero after inverting  $p$ .  $\square$

**Remark 5.2.** For  $d \geq 4$ , this result is due to Arone, Lambrechts, Turchin and Volić (see [1]). The case  $d = 3$  does not appear in *loc. cit.* and, to the best of our knowledge, does not appear elsewhere in the literature. The reason seems to be that the relative formality of the map of operads  $E_1 \rightarrow E_3$  was not known when [1] was written.

**Corollary 5.3.** *Let  $p$  be a prime. For  $n \leq (p-1)(d-2) + 3$ , the spectral sequence associated to  $T_n(\mathbb{Z}_p)$ , computing*

$$\pi_* T_n \text{emb}_c(\mathbb{R}^1, \mathbb{R}^d) \otimes \mathbb{Z}_p,$$

*collapses at the  $E^2$ -page for  $(-s, t)$  satisfying  $t < 2p - 2 + (s-1)(d-1)$ .*

*Proof.* Let  $T(s) := 2p - 2 + (s-1)(d-1)$ . Let  $A$  be the region of the second-quadrant consisting of those  $(-s, t)$  such that  $t < T(s)$ . By theorem 4.1, the first possibly non-trivial differential in the region  $A$  is

$$d^R : E_{-s,t}^R \rightarrow E_{-s-R,t+R-1}^R.$$

with  $R = (p-1)(d-2) + 1$ . Since the first non-zero column is  $s = 3$ , the first possibly non-trivial differentials land in  $E_{-3-R,t+R-1}^R$ . But the groups  $E_{-s,*}^1$  are zero whenever  $s > n$ . So  $E_{-3-R,t+R-1}^R$  is zero, and so are the target groups of higher differentials from the region  $A$ . Therefore, the spectral sequence collapses in the region  $A$  whenever  $n < R + 3$ .  $\square$

**Remark 5.4.** Note that the corollary has nothing to offer at the prime 2.

**Remark 5.5.** Theorem 4.1 has something to say about the homotopy groups of  $\pi_*(T_n)$  even outside of the range of the previous corollary. For instance, let us consider  $\pi_0(T_n) \otimes \mathbb{Z}_p$  in the case of knots in  $\mathbb{R}^3$ . This abelian group has filtration whose associated graded is  $\bigoplus_{s \leq n} E_{-s,s}^\infty(T_n(\mathbb{Z}_p))$ . The corollary tells us that we have isomorphisms

$$E_{-s,s}^2 \cong E_{-s,s}^\infty$$

in the range  $s \leq p + 2$ . For  $p + 3 \leq s \leq 2p + 1$ , the only differential that can hit  $E_{-s,s}^2$  is  $d^p$ , therefore, we have an isomorphism

$$E_{-s,s}^\infty \cong E_{-s,s}^{p+1} \cong E_{-s,s}^2 / \text{Im}(d^p)$$

For  $2p + 2 \leq s \leq 3p$ , the group  $E_{-s,s}^\infty$  will be the quotient of  $E_{-s,s}^2$  by the image of  $d^p$  and  $d^{2p-1}$ . This pattern continues.

In the remainder of this section, we prove some results about extensions leading to the proof of Theorem C from the introduction. Let us fix a unit  $u \in \mathbb{Z}_p$  such that the residue of  $u$  modulo  $p$  is a generator of  $\mathbb{F}_p^\times$ . For  $M$  a  $\mathbb{Z}_p$ -module and  $i$  an integer, we denote by  $M(i)$  the  $\mathbb{Z}_p[t, t^{-1}]$ -module  $M$  where  $t$  acts as multiplication by  $u^i$ . When  $i = 0$ , we simply write  $M$  for  $M(0)$ .

**Proposition 5.6.** *Let  $M$  be a  $\mathbb{Z}_p$ -module of finite type. Let  $i$  and  $j$  be two integers such that  $(p-1)$  does not divide  $i - j$ . Then the group  $\text{Ext}_{\mathbb{Z}_p[t, t^{-1}]}^1(M(i), N(j))$  is zero.*

*Proof.* Since  $M$  can be written as finite direct sums of copies of  $\mathbb{Z}_p$  and  $\mathbb{Z}/p^n$ , without loss of generality, we may assume that  $M$  is either  $\mathbb{Z}_p$  or  $\mathbb{Z}/p^n$ . The short exact sequence

$$0 \rightarrow \mathbb{Z}/p^n \rightarrow \mathbb{Z}/p^{n-1} \rightarrow \mathbb{Z}/p \rightarrow 0$$

induces an exact sequence

$$\begin{aligned} \mathrm{Ext}_{\mathbb{Z}_p[t,t^{-1}]}^1(\mathbb{Z}/p(i), N(j)) &\rightarrow \mathrm{Ext}_{\mathbb{Z}_p[t,t^{-1}]}^1(\mathbb{Z}/p^{n+1}(i), N(j)) \\ &\rightarrow \mathrm{Ext}_{\mathbb{Z}_p[t,t^{-1}]}^1(\mathbb{Z}/p^n(i), N(j)) \end{aligned}$$

Therefore, by induction on  $n$ , we can reduce the case  $M = \mathbb{Z}/p^n$  to the case  $M = \mathbb{Z}/p$ . The short exact sequence

$$0 \rightarrow \mathbb{Z}_p \xrightarrow{p} \mathbb{Z}_p \rightarrow \mathbb{Z}/p \rightarrow 0$$

induces an exact sequence

$$\mathrm{Hom}_{\mathbb{Z}_p[t,t^{-1}]}(\mathbb{Z}_p(i), N(j)) \rightarrow \mathrm{Ext}_{\mathbb{Z}_p[t,t^{-1}]}^1(\mathbb{Z}/p(i), N(j)) \rightarrow \mathrm{Ext}_{\mathbb{Z}_p[t,t^{-1}]}^1(\mathbb{Z}_p(i), N(j))$$

therefore, the case  $M = \mathbb{Z}/p$  follows from the vanishing of  $\mathrm{Hom}_{\mathbb{Z}_p[t,t^{-1}]}(\mathbb{Z}_p(i), N(j))$  and of  $\mathrm{Ext}_{\mathbb{Z}_p[t,t^{-1}]}^1(\mathbb{Z}_p(i), N(j))$ . The vanishing of the Hom-group is similar to the proof of Proposition 2.2. In order to compute the Ext-group, we use the following projective resolution of  $\mathbb{Z}_p(i)$

$$0 \rightarrow \mathbb{Z}_p[t, t^{-1}] \xrightarrow{\cdot(t-u^i)} \mathbb{Z}_p[t, t^{-1}] \rightarrow \mathbb{Z}_p \rightarrow 0.$$

We obtain an exact sequence

$$N \rightarrow N \rightarrow \mathrm{Ext}_{\mathbb{Z}_p[t,t^{-1}]}^1(\mathbb{Z}_p(i), N(j)) \rightarrow 0$$

where the first map is multiplication by  $u^j - u^i$ . Since  $j - i$  is not a multiple of  $(p - 1)$ ,  $u^j - u^i$  is a unit of  $\mathbb{Z}_p$  and the result follows.  $\square$

**Proposition 5.7.** *Let  $M$  be a  $\mathbb{Z}_p$ -module of finite type equipped with a  $GT_p$ -action. Let  $n$  and  $m$  be two integers with  $0 \leq n - m < p - 1$ . Assume that  $M$  sits at the top of a  $GT_p$ -equivariant tower*

$$M = M_n \xrightarrow{p_n} M_{n-1} \xrightarrow{p_{n-1}} \dots \rightarrow M_m \xrightarrow{p_m} 0$$

where each map  $p_k$  is surjective. Assume further that, for each  $k$ , the  $GT_p$ -action on the kernel  $F_k$  of  $p_k$  is cyclotomic of weight  $k$ . Then there is a non-canonical isomorphism of  $\mathbb{Z}_p$ -modules

$$M \cong F_n \oplus \dots \oplus F_m.$$

*Proof.* It suffices to prove that, for each  $k$ , the short exact sequence

$$0 \rightarrow F_k \rightarrow M_k \rightarrow M_{k-1} \rightarrow 0$$

is split. The obstruction lies in the Ext-group  $\mathrm{Ext}_{\mathbb{Z}_p}^1(M_{k-1}, F_k)$ . Since the short exact sequence is  $GT_p$ -equivariant, the obstruction lies in the image of the obvious map

$$\mathrm{Ext}_{\mathbb{Z}_p[GT]}^1(M_{k-1}, F_k) \rightarrow \mathrm{Ext}_{\mathbb{Z}_p}^1(M_{k-1}, F_k).$$

Pick an element  $u$  of  $\mathbb{Z}_p^\times$  such that the residue of  $u$  modulo  $p$  is a generator of  $\mathbb{F}_p^\times$ . Pick a lift  $t$  of  $u$  in  $GT_p$ . This choice induces a homomorphism of  $\mathbb{Z}_p$ -algebras

$$\mathbb{Z}_p[t, t^{-1}] \rightarrow \mathbb{Z}_p[GT]$$

and we deduce that the obstruction lies in the image of the obvious map

$$\mathrm{Ext}_{\mathbb{Z}_p[t,t^{-1}]}^1(M_{k-1}, F_k) \rightarrow \mathrm{Ext}_{\mathbb{Z}_p}^1(M_{k-1}, F_k)$$



Hence it is enough to show that the group  $\text{Ext}_{\mathbb{Z}_p[t,t^{-1}]}^1(M_{k-1}, F_k)$  is zero. We will in fact prove that the group  $\text{Ext}_{\mathbb{Z}_p[t,t^{-1}]}^1(M_i, F_k) = 0$  for any  $m \leq i < k$ . We prove this by induction on  $i$ . For  $i = m$ , we have

$$\text{Ext}_{\mathbb{Z}_p[t,t^{-1}]}^1(M_m, F_k) = 0$$

by the previous proposition. If the statement is true for  $i$ , then, we use the long exact sequence induced by the short exact sequence

$$0 \rightarrow F_{i+1} \rightarrow M_{i+1} \rightarrow M_i \rightarrow 0$$

to conclude.  $\square$

Using this proposition, we can prove Theorem C from the introduction.

*Proof of Theorem C.* The analogous statement with  $\mathbb{Z}_p$  replaced by  $\mathbb{Z}_{(p)}$  follows from Corollary 5.3 and the previous proposition. If  $M$  and  $N$  are finitely generated  $\mathbb{Z}_{(p)}$ -modules, the canonical map

$$\text{Ext}(M, N) \otimes \mathbb{Z}_p \rightarrow \text{Ext}_{\mathbb{Z}_p}(M \otimes \mathbb{Z}_p, N \otimes \mathbb{Z}_p)$$

is an isomorphism. This can be reduced to the case where  $M$  and  $N$  are either  $\mathbb{Z}_{(p)}$  or  $\mathbb{Z}/p^k$  in which case this is an easy computation. It follows that if an extension of  $\mathbb{Z}_{(p)}$ -modules splits over  $\mathbb{Z}_p$ , it also splits over  $\mathbb{Z}_{(p)}$ .  $\square$

## 6. UNIVERSALITY

We now come to the proof of theorem A from the introduction. To lighten notations, we denote the space of knots  $\text{emb}_c(\mathbb{R}^1, \mathbb{R}^3)$  by  $K$  in this section, as we did in the introduction. Given two knots  $f, g$ , we write  $f \sim_{n-1} g$  if  $f$  and  $g$  share the same type  $n-1$  invariants. This defines an equivalence relation on  $\pi_0 K$ ; the set of equivalence classes is denoted  $\pi_0(K)/\sim_{n-1}$ . It is not hard to see that the operation of concatenation makes this set into a commutative monoid. This commutative monoid is in fact a finitely generated abelian group as shown in [17].

**Theorem 6.1.** *Let  $R$  be a commutative ring that is torsion-free (e.g.  $\mathbb{Z}$ ,  $\mathbb{Q}$ ,  $\mathbb{Z}_p$ ,  $\mathbb{Z}_{(p)}$ ). The evaluation map*

$$e_n : \pi_0 K \rightarrow \pi_0 T_n K \otimes R$$

*is a universal Vassiliev invariant of degree  $n-1$  over  $R$  if the canonical map*

$$E_{-k,k}^2 \otimes R \rightarrow E_{-k,k}^\infty \otimes R$$

*is an isomorphism for all  $k \leq n$ .*

*Proof.* In [8], and [21], it was shown that  $e_n$  is well-defined on equivalence classes and as such factors through  $\bar{e}_n : (\pi_0(K)/\sim_{n-1}) \rightarrow \pi_0 T_n K$ . The statement of the theorem is equivalent to the statement that  $\bar{e}_n \otimes R$  is an isomorphism.

We argue inductively, using the commutative square of group homomorphisms

$$\begin{array}{ccc} \pi_0(K)/\sim_{n-1} & \xrightarrow{\bar{e}_n} & \pi_0 T_n K \\ \downarrow & & \downarrow \\ \pi_0(K)/\sim_{n-2} & \xrightarrow{\bar{e}_{n-1}} & \pi_0 T_{n-1} K \end{array}$$

and compare the map between vertical kernels

$$\Phi_{n-1} \rightarrow E_{-n,n}^\infty$$

where  $\Phi_{n-1}$  denotes the kernel of the left vertical map. In [11], Conant-Teichner construct a surjective homomorphism  $R_n : E_{-n,n}^2 \rightarrow \Phi_{n-1}$  and in [20], Kosanović shows that the composition

$$E_{-n,n}^2 \rightarrow \Phi_{n-1} \rightarrow E_{-n,n}^\infty$$

agrees with the canonical map. Since the composition is an isomorphism by hypothesis, we then have that both maps are isomorphisms. It follows by an application of the five lemma that  $\bar{e}_n \otimes R$  is an isomorphism if and only if the homomorphism  $E_{-n,n}^2 \otimes R \rightarrow E_{-n,n}^\infty \otimes R$  is an isomorphism, as claimed.  $\square$

*Proof of Theorem A.* The first part follows from Theorem 6.1 in conjunction with Theorem B. The second part is subsumed by Theorem C.  $\square$

## 7. THE HOMOLOGY GOODWILLIE-WEISS SPECTRAL SEQUENCE

The Goodwillie-Weiss tower for  $\overline{\text{emb}}_c(\mathbb{R}, \mathbb{R}^d)$ , the homotopy fiber of the inclusion

$$\text{emb}_c(\mathbb{R}, \mathbb{R}^d) \rightarrow \text{imm}_c(\mathbb{R}, \mathbb{R}^d),$$

can also be described as follows. Consider the category  $\mathcal{O}_k$  whose objects are open proper subsets  $T$  of  $\mathbb{R}$  containing  $(-\infty, 0] \cup [1, +\infty)$  and having at most  $k+2$  connected components. A morphism  $T \rightarrow R$  is an isotopy connecting  $T \subset \mathbb{R}$  to a subset of  $R \subset \mathbb{R}$ , and which fixes  $(-\infty, 0] \cup [1, +\infty)$  pointwise. Then

$$T_k \overline{\text{emb}}_c(\mathbb{R}^1, \mathbb{R}^d) = \text{holim}_{T \in \mathcal{O}_k^{\text{op}}} \overline{\text{emb}}_c(T, \mathbb{R}^d).$$

The Goodwillie-Weiss tower agrees with the tower associated to a cosimplicial space, the one coming from the usual filtration of  $\Delta$  by the subcategories  $\Delta_{\leq k}$  spanned by the objects  $[n]$  satisfying  $n \leq k$ . This was first proved by Sinha [25]. Roughly, the relation comes from the equivalence between the topological category  $\mathcal{O}_k$  and the opposite of  $\Delta_{\leq k}$ . It follows that the functor  $\overline{\text{emb}}_c(-, \mathbb{R}^d)$  defines a cosimplicial space whose homotopy limit is the limit of the Goodwillie-Weiss tower. In this section, we review an operadic construction of this cosimplicial space and use it to study the associated homology Bousfield-Kan spectral sequence.

Recall that a multiplicative operad is a non-symmetric topological or simplicial operad  $P$  together with the data of a map  $\alpha : A \rightarrow P$  where  $A$  denotes the non-symmetric associative operad. From such data, one can construct a cosimplicial space  $X^\bullet$  as we now recall (see [25, Definition 2.17]). We denote by  $u$  the image in  $P(0)$  of the unique point in  $A(0)$  and by  $m$  the image in  $P(2)$  of the unique point in  $A(2)$ .

In degree  $q$ , our cosimplicial space will be given by  $X^q = P(q)$ . The cofaces  $d^i : P(q) \rightarrow P(q+1)$  are as follows. The two outer cofaces  $d^0$  and  $d^{q+1}$  take  $x \in P(q)$  to  $m \circ_1 x$  and  $m \circ_2 x$  respectively. The inner coface  $d^i$  with  $i \in \{1, \dots, q\}$  takes  $x$  to  $x \circ_i m$ . The codegeneracy  $s^i : P(q) \rightarrow P(q-1)$  takes  $x \in P(q)$  to  $x \circ_i u$ .

In what follows, we will start with a map  $\tilde{A} \rightarrow P$  where  $\tilde{A}$  is not quite the associative operad but is merely weakly equivalent to it. The following proposition will be useful in that situation. We denote by  $\mathbf{Op}_{n_s}$  the category of non-symmetric operads in simplicial sets.

**Proposition 7.1.** *Let  $\tilde{A}$  be a non-symmetric operad and let  $w : \tilde{A} \rightarrow A$  be a map of non-symmetric operads. Consider the adjunction*

$$w_! : \mathbf{Op}_{ns}^{\tilde{A}/} \rightleftarrows \mathbf{Op}_{ns}^{A/} : w^*$$

where  $w^*$  is precomposition by  $w$  and  $w_!$  is its left adjoint. Then, if  $w$  is a weak equivalence, the adjunction is a Quillen equivalence.

*Proof.* First, we easily verify that the left adjoint sends  $\tilde{A} \rightarrow P$  to the bottom map in the following pushout square.

$$\begin{array}{ccc} \tilde{A} & \longrightarrow & P \\ \downarrow & & \downarrow \\ A & \longrightarrow & Q \end{array}$$

The result is then completely standard once we know that the model category  $\mathbf{Op}_{ns}$  is left proper (see [23, Corollary 1.12]).  $\square$

Fix a linear inclusion  $\mathbb{R} \rightarrow \mathbb{R}^d$ . This induces a map of non-symmetric operads  $E_1^{ns} \rightarrow E_d$  where  $E_1^{ns}$  is the non-symmetric little 1-disks operad. The unique map  $w : E_1^{ns} \rightarrow A$  being a weak equivalence, we can apply the derived functor of  $w_!$  (using the notation of Proposition 7.1) to it and get a map  $A \rightarrow P$  for some non-symmetric operad  $P$ . Moreover, by Proposition 7.1 this map is weakly equivalent to  $E_1^{ns} \rightarrow E_d$  in the arrow category of  $\mathbf{Op}^{ns}$ . The cosimplicial space associated to this map is what we will denote by  $K^\bullet$ . It is identified with the cosimplicial space mentioned in the beginning of the section.

We can do the exact same construction but starting from the map  $L_p E_1 \rightarrow L_p E_d$  (noticing that  $L_p E_1 \simeq E_1$ ) and we obtain a cosimplicial space that we denote by  $L_p K^\bullet$ . Since the map  $L_p E_1 \rightarrow L_p E_d$  is  $GT_p$ -equivariant (Theorem 2.3), the cosimplicial space  $L_p K_d^\bullet$  gets a  $GT_p$ -action.

Applying the functor  $C_*(-, A)$  to the cosimplicial space  $K^\bullet$ , we get a bicomplex and hence a spectral sequence  $E_{*,*}^*(A)$  with

$$E_{p,q}^1(A) = H_q(K_d^{-p}, A)$$

and whose  $d_1$  is the alternating sum of the coface maps. Our main theorem about this spectral sequence is the following (Theorem D in the introduction).

**Theorem 7.2.** *Let  $p$  be a prime number. The only possibly non-trivial differentials in the spectral sequence  $E^*(\mathbb{Z}_{(p)})$  are  $d_{1+n(d-1)(p-1)}$  for  $n \geq 0$ .*

*Proof.* As in the proof of Theorem B, it suffices to prove the theorem for the spectral sequence  $E^*(\mathbb{Z}_p)$ . As we said above, we have a  $GT_p$ -action on the cosimplicial space  $L_p K_d^\bullet$ . Using Proposition 1.5, we obtain an action on  $C^*(X, \mathbb{Z}_p)$  and hence on the spectral sequence  $E^*(\mathbb{Z}_p)$ . Moreover, the action on  $E_{*,(d-1)k}^*$  is cyclotomic of weight  $k$ . Indeed, this is true on the  $E^1$  page by Proposition 2.4 and therefore on any page by Proposition 2.1. The result now follows from Proposition 2.2.  $\square$

In particular, we recover the following theorem.

**Corollary 7.3** (Lambrechts-Turchin-Volić, [22]). *The spectral sequence  $E^*(\mathbb{Q})$  collapses at the second page.*

*Proof.* The proof is analogous to the proof of Corollary 5.1.  $\square$

## REFERENCES

- [1] ARONE, G., LAMBRECHTS, P., TURCHIN, V., AND VOLIĆ, I. Coformality and rational homotopy groups of spaces of long knots. *Mathematical Research Letters* 15, 1 (2008), 1–15.
- [2] BAR-NATAN, D. On the Vassiliev knot invariants. *Topology* 34 (1995), 423–472.
- [3] BOAVIDA DE BRITO, P., AND HOREL, G. On the formality of the little disks operad in positive characteristic. arXiv:1903.09191, 2019.
- [4] BOAVIDA DE BRITO, P., AND WEISS, M. Spaces of smooth embeddings and configuration categories. *Journal of Topology* 11, 1 (2018), 65–143.
- [5] BOAVIDA DE BRITO, P., AND WEISS, M. S. The configuration category of a product. *Proc. Amer. Math. Soc.* 146, 10 (2018), 4497–4512.
- [6] BOUSFIELD, A. K. The localization of spaces with respect to homology. *Topology* 14 (1975), 133–150.
- [7] BOUSFIELD, A. K., AND KAN, D. M. *Homotopy limits, completions and localizations*. Lecture Notes in Mathematics, Vol. 304. Springer-Verlag, Berlin-New York, 1972.
- [8] BUDNEY, R., CONANT, J., KOYTSCHEFF, R., AND SINHA, D. Embedding calculus knot invariants are of finite type. *Algebraic and Geometric Topology* 17, 3 (2017), 1701–1742.
- [9] BUDNEY, R., CONANT, J., SCANNELL, K. P., AND SINHA, D. New perspectives on self-linking. *Advances in Mathematics* 191, 1 (2005), 78 – 113.
- [10] CONANT, J. Homotopy approximations to the space of knots, Feynman diagrams, and a conjecture of Scannell and Sinha. *American Journal of Mathematics* 130, 2 (2008), 341–357.
- [11] CONANT, J., AND TEICHNER, P. Grope cobordism and Feynman diagrams. *Math. Ann.* 328, 1-2 (2004), 135–171.
- [12] DRINFEL'D, V. G. On quasitriangular quasi-Hopf algebras and on a group that is closely connected with  $\text{Gal}(\overline{\mathbf{Q}}/\mathbf{Q})$ . *Algebra i Analiz* 2, 4 (1990), 149–181.
- [13] DWYER, W., AND HESS, K. Long knots and maps between operads. *Geometry & Topology* 16, 2 (2012), 919–955.
- [14] FURUSHO, H. Galois action on knots I: Action of the absolute Galois group. *Quantum Topology* 8, 2 (2017), 295–360.
- [15] GOODWILLIE, T. G., AND WEISS, M. Embeddings from the point of view of immersion theory. II. *Geom. Topol.* 3 (1999), 103–118.
- [16] GÖPPL, F. A spectral sequence for spaces of maps between operads. *arXiv e-prints* (Oct 2018), arXiv:1810.05589.
- [17] GUSAROV, M. On  $n$ -equivalence of knots and invariants of finite degree. In *Topology of manifolds and varieties*, vol. 18 of *Adv. Soviet Math.* Amer. Math. Soc., Providence, RI, 1994, pp. 173–192.
- [18] HOREL, G. Groupe de galois et espace des noeuds. In *SMF 2018 : Congrès de la SMF*, vol. 33 of *Sémin. Congr. Soc. Math. France*, Paris, 2019, pp. 273–282.
- [19] KASSEL, C., AND TURAEV, V. Chord diagram invariants of tangles and graphs. *Duke Math. J.* 92, 3 (1998), 497–552.
- [20] KOSANOVIĆ, D. A geometric approach to the embedding calculus knot invariants. December 2019.
- [21] KOSANOVIĆ, D., SHI, Y., AND TEICHNER, P. Space of gropes and the embedding calculus. *In preparation*.
- [22] LAMBRECHTS, P., TURCHIN, V., AND VOLIĆ, I. The rational homology of spaces of long knots in codimension  $> 2$ . *Geometry & Topology* 14, 4 (2010), 2151–2187.
- [23] MURO, F. Homotopy theory of non-symmetric operads, II: Change of base category and left properness. *Algebraic & Geometric Topology* 14, 1 (2014), 229–281.
- [24] SHI, Y. Vassiliev invariants via manifold calculus. Master's thesis, University of Bonn, 2018.
- [25] SINHA, D. Operads and knot spaces. *Journal of the American Mathematical Society* 19, 2 (2006), 461–486.
- [26] TURCHIN, V. Delooping totalization of a multiplicative operad. *J. Homotopy Relat. Struct.* 9, 2 (2014), 349–418.
- [27] VASIL'EV, V. A. *Complements of discriminants of smooth maps: topology and applications. Transl. from the Russian by B. Goldfarb. Transl. ed. by S. Gelfand. Rev. ed.*, vol. 98. Providence, RI: American Mathematical Society, 1994.
- [28] VOLIC, I. Finite type knot invariants and the calculus of functors. *Compositio Mathematica* 142, 01 (2006), 222–250.

DEPT. OF MATHEMATICS, IST, UNIV. OF LISBON, AV. ROVISCO PAIS, LISBOA, PORTUGAL  
*Email address:* `pedrobrito@tecnico.ulisboa.pt`

UNIVERSITÉ SORBONNE PARIS NORD, LAGA, CNRS, UMR 7539, F-93430, VILLETANEUSE,  
FRANCE

ÉCOLE NORMALE SUPÉRIEURE, DMA, CNRS, UMR 8553, 45 RUE D'ULM, 75230 PARIS CEDEX  
05, FRANCE  
*Email address:* `horel@math.univ-paris13.fr`